

Available online at www.sciencedirect.com

SciVerse ScienceDirect

www.nrjournal.com

Communication

Curcumin ingestion and exercise training improve vascular endothelial function in postmenopausal women

Nobuhiko Akazawa^a, Youngju Choi^b, Asako Miyaki^a, Yoko Tanabe^a, Jun Sugawara^c,
Ryuichi Ajisaka^b, Seiji Maeda^{b,*}

^a Graduate School of Comprehensive Human Science, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki, Japan

^b Faculty of Health and Sport Science, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki, Japan

^c Human Technology Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), 1-1-1 Higashi, Tsukuba, Ibaraki, Japan

ARTICLE INFO

Article history:

Received 11 April 2012

Revised 19 August 2012

Accepted 12 September 2012

Keywords:

Flow-mediated dilation

Turmeric

Physical activity

Lifestyle modification

Menopause

Women

ABSTRACT

Vascular endothelial function declines with aging and is associated with an increased risk of cardiovascular disease. Lifestyle modification, particularly aerobic exercise and dietary adjustment, has a favorable effect on vascular aging. Curcumin is a major component of turmeric with known anti-inflammatory and anti-oxidative effects. We investigated the effects of curcumin ingestion and aerobic exercise training on flow-mediated dilation as an indicator of endothelial function in postmenopausal women. A total of 32 postmenopausal women were assigned to 3 groups: control, exercise, and curcumin groups. The curcumin group ingested curcumin orally for 8 weeks. The exercise group underwent moderate aerobic exercise training for 8 weeks. Before and after each intervention, flow-mediated dilation was measured. No difference in baseline flow-mediated dilation or other key dependent variables were detected among the groups. Flow-mediated dilation increased significantly and equally in the curcumin and exercise groups, whereas no changes were observed in the control group. Our results indicated that curcumin ingestion and aerobic exercise training can increase flow-mediated dilation in postmenopausal women, suggesting that both can potentially improve the age-related decline in endothelial function.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Arterial endothelial dysfunction has been associated with cardiovascular morbidity and mortality [1]. Flow-mediated dilation (FMD), which is an index of endothelial function [2], is progressively impaired with age [3,4]. In women, the age-associated decline in endothelial function is enhanced

during menopause because of the estrogen deficiency [5,6], and postmenopausal women are at a higher risk of cardiovascular disease [7]. Thus, slowing the decline of endothelial function in postmenopausal women could have potential health benefits. We and other groups have reported that aerobic exercise training enhances endothelial function [8–10]. Lifestyle modification is a desirable way to prevent or

Abbreviations: FMD, flow-mediated dilation; VO_{2peak} , peak oxygen consumption; HDL cholesterol, high-density lipoprotein cholesterol; LDL cholesterol, low-density lipoprotein cholesterol; TNF- α , tumor necrosis factor-alpha.

* Corresponding author. Tel.: +81 29 853 2683; fax: +81 29 853 2986.

E-mail address: maeda@taiiku.tsukuba.ac.jp (S. Maeda).

0271-5317/\$ – see front matter © 2012 Elsevier Inc. All rights reserved.

<http://dx.doi.org/10.1016/j.nutres.2012.09.002>

treat endothelial dysfunction without the need for pharmaceutical intervention.

Vascular inflammation and oxidative stress play important roles in development of vascular endothelial dysfunction and cardiovascular disease [11]. Previous studies have demonstrated that increased production of pro-inflammatory cytokines and reactive oxygen species in the aging vessel results in endothelial dysfunction [12,13]. Therefore, following an anti-inflammatory or anti-oxidative diet may help counteract the effects of aging on blood vessel. Curcumin, a polyphenol molecule extracted from turmeric, is a commonly used spice and a yellow pigment. Curcumin regulate biochemical and molecular pathway by modulating several molecular targets including transcription factors, cytokines, enzymes, and genes regulating cell proliferation and apoptosis [14–16]. In addition to the anti-inflammatory and anti-oxidative effects of curcumin [17,18], it has been associated with the protection against cardiovascular disease [19]. However, the effect of curcumin on endothelial function remains unclear.

We hypothesized that, similar to exercise, curcumin ingestion could improve endothelial function. The objective of the present study was to determine the effect of curcumin ingestion and the effect of exercise training on endothelial function. We therefore investigated endothelial function as measured by the FMD in postmenopausal women before and after 8 weeks of curcumin ingestion or exercise training intervention.

2. Methods and materials

2.1. Subjects

A total of 32 healthy, sedentary postmenopausal women (amenorrhea for at least 2 years) participated in the study. Subjects were assigned to one of the following intervention groups: control group ($n = 10$), curcumin group ($n = 11$), and exercise training group ($n = 11$). Subjects were nonsmokers, nonobese, and free of cardiovascular disease as assessed by medical history. None of the subjects were taking cardiovascular-acting medications or hormone replacement therapy. All potential risks and associated with the study were explained to the subjects, and they gave their written informed consent for participation in the study. All procedures were reviewed and approved by the ethical committee of the University of Tsukuba.

2.2. Experimental protocol

All experiments were performed in the morning after a 12-h overnight fast. Subjects abstained from alcohol and caffeine for at least 12 h and did not exercise for at least 24 h before beginning the experiment to avoid the potential acute effects of exercise. Measurements were taken in a quiet, temperature-controlled room (24–26°C). After a resting period of at least 20 min, FMD, arterial blood pressure, and blood biochemistry were determined. After these measurements, peak oxygen consumption (VO_{2peak}) was measured during incremental cycle ergometer exercise.

2.3. Curcumin ingestion

Subjects in the curcumin group ingested 6 pills (150 mg total) of curcumin per day [20,21], which supplies 25 mg of highly absorptive curcumin dispersed with colloidal nanoparticles (Theracurmin; Theravalues Corporation, Tokyo, Japan) [22]. Supplementary curcumin was administered orally for 8 weeks. All subjects were instructed not to alter their dietary habits during the intervention period.

2.4. Exercise training

Subjects in the exercise group underwent aerobic exercise training more than 3 days per week (2–3 supervised sessions and additional home-based training) for 8 weeks [23]. Initially, subjects performed cycling and walking 30 min/d at a relatively low intensity (60% of their individually determined maximal heart rate). As their exercise tolerance improved, the intensity and duration of aerobic exercise were increased to 40 to 60 min/d at an intensity of 70% to 75% of the maximal heart rate. Subjects in the control and curcumin groups were instructed not to change their level of physical activity.

2.5. Measurements

2.5.1. FMD

Brachial artery FMD was assessed noninvasively by using an ultrasound system (UNEXEF18G; Unex, Nagoya, Japan) as previously described [24]. Briefly, high-resolution ultrasound with a 10 MHz linear array transducer was used to obtain a longitudinal image of the right brachial artery in the cubital region at the baseline and then continuously from 30 seconds prior to 2 minutes or more after the release of suprasystolic pressure (50 mm Hg above systolic blood pressure) maintained for 5 minutes of the right forearm. The diameter at the same point of the artery was monitored continuously, and the maximal dilatation after deflation was recorded. FMD was calculated as the percentage change in brachial artery diameter in response to the forearm reactive hyperemic stimulus. FMD was calculated as follows: $\text{Maximal diameter} - \text{baseline diameter} \times 100 / \text{baseline diameter}$

2.5.2. Arterial blood pressure

Arterial blood pressure and heart rate at rest were determined in the supine position using an automated device (form PWV/ABI, Colin Medical Technology, Komaki, Japan) [23].

2.5.3. Blood chemistry

A blood sample was collected from the antecubital vein after overnight fasting. Serum of cholesterol and triglyceride and plasma of glucose were determined using standard enzymatic techniques [25].

2.5.4. Peak oxygen consumption

VO_{2peak} was measured during incremental cycle ergometer exercise by using online computer-assisted circuit spirometry (AE280, Minato Medical Science, Osaka, Japan) as we previously reported [23]. All subjects underwent an incremental exercise test (2 minutes at 40 W, followed by 20 W increases every 2 minutes) until volitional exhaustion. VO_{2peak} was defined as the highest VO_2 recorded during the test. Heart rate and rating of perceived exertion were recorded throughout the exercise.

Table 1 – Baseline characteristics of subjects

	Control (n = 10)		Exercise (n = 11)		Curcumin (n = 11)		P		
	Before	After	Before	After	Before	After	Group	Time	Time × group
Age, years	64 ± 6		59 ± 5		60 ± 6		NS		
Height, cm	154 ± 6		154 ± 4		155 ± 8		NS		
Weight, kg	51 ± 2	51 ± 2	54 ± 2	54 ± 2	54 ± 3	54 ± 3	NS	NS	NS
BMI, kg/m ²	21.5 ± 1.0	21.6 ± 1.0	22.7 ± 1.0	22.6 ± 1.0	22.5 ± 1.0	22.7 ± 1.0	NS	NS	NS
Total cholesterol, mmol/L	6.1 ± 0.5	6.3 ± 0.5	5.4 ± 0.6	5.7 ± 0.6	6.3 ± 0.7	6.6 ± 0.5	NS	0.005	NS
HDL cholesterol, mmol/L	1.7 ± 0.4	1.7 ± 0.3	1.7 ± 0.3	1.9 ± 0.3*	1.6 ± 0.5	1.7 ± 0.5	NS	0.010	NS
LDL cholesterol, mmol/L	3.8 ± 0.6	3.9 ± 0.4	3.2 ± 0.4	3.5 ± 0.6	3.8 ± 0.7	4.0 ± 0.7	NS	0.027	NS
Triglyceride, mmol/L	1.4 ± 0.6	1.3 ± 0.4	1.0 ± 0.6	0.9 ± 0.4	1.9 ± 0.9	1.7 ± 0.1	NS	NS	NS
Glucose, mmol/L	5.3 ± 0.4	5.3 ± 0.4	4.9 ± 0.4	4.9 ± 0.3	5.1 ± 1.1	5.4 ± 1.6	NS	NS	NS
VO _{2peak} , mL/kg/min	22.9 ± 1.4	22.5 ± 1.1	25.3 ± 1.2	27.3 ± 1.2*	21.9 ± 0.7	21.9 ± 0.8	NS	NS	0.024

Values are means ± SD. BMI; body mass index.

* P < .05 vs. before intervention. A main time and group effect overall by repeated-measures analysis of variance.

2.6. Statistical analyses

The effects of each intervention on the outcomes measured were determined by repeated measures analysis of variance. If a significant effect was detected, specific mean comparisons were performed to determine the significance of each intervention. In the case of a significant F value, a post hoc test (Bonferroni test) was used to identify significant differences among mean values. All data are reported as means ± SD. Statistical significance was set a priori at P < .05 for all comparisons.

3. Results

The compliance/adherence to curcumin ingestion in the curcumin group was 97.6% ± 2.8%. No adverse effects of curcumin were reported. In the exercise training group, the average frequency and duration of the exercise training was 4.5 ± 1.4 d/wk and 48 ± 23 min/d. There were no dropouts in either group.

Table 1 shows the baseline characteristics of the study participants. There were no group differences in age, height, body weight, body mass index, total cholesterol, high-density lipoprotein (HDL) cholesterol, low-density lipoprotein (LDL) cholesterol, triglyceride, glucose, or VO_{2peak} between the groups at the start of the study. HDL cholesterol levels increased significantly in the exercise group (P < .05). There were no significant changes in body weight, body mass index, total cholesterol, LDL cholesterol, triglyceride and glucose levels in any of the groups. VO_{2peak} in the exercise group increased significantly after the intervention (P < .05).

3.1. Heart rate and blood pressure

As shown in Table 2, no statistically significant differences in the baseline hemodynamic parameters at rest were detected among the groups before the intervention. After 8 weeks of intervention, systolic blood pressure decreased in the exercise and curcumin groups (P < .05). Heart rate and diastolic blood pressure did not change in any of the groups.

3.2. FMD

No significant difference in the baseline FMD was detected among the groups (Fig. 1). After 8 weeks of intervention, FMD increased significantly in the exercise and curcumin groups, whereas no significant change in FMD was detected in the control group. The change in FMD was significantly greater in the exercise and the curcumin groups than in the control group (Fig. 2). The differences in FMD changes between the exercise and curcumin groups were not statistically significant (Fig. 2).

4. Discussion

The present study showed that regular ingestion of curcumin or regular aerobic exercise training significantly improved endothelial function. The magnitude of improvement in endothelial function to the same extent, suggesting that curcumin may prevent the age-associated decline in endothelial function in postmenopausal women.

Table 2 – Hemodynamics parameter before and after intervention

	Control (n = 10)		Exercise (n = 11)		Curcumin (n = 11)		P		
	Before	After	Before	After	Before	After	Group	Time	Time × group
Heart rate, beats/min	62 ± 6	64 ± 11	61 ± 6	57 ± 5	61 ± 8	59 ± 7	NS	NS	NS
SBP, mmHg	112 ± 12	113 ± 11	112 ± 10	107 ± 10*	122 ± 17	116 ± 15*	NS	0.003	0.025
DBP mmHg	69 ± 5	69 ± 6	71 ± 8	69 ± 9	72 ± 12	69 ± 11	NS	NS	NS

Values are means ± SD. SBP; systolic blood pressure, DBP; diastolic blood pressure.

* P < .05 vs. before intervention. A main time and group effect overall by repeated-measures analysis of variance.




Fig. 1 – FMD before and after intervention. Data are expressed as means \pm SD. * $P < .05$ before vs after intervention.

Curcumin, a polyphenolic derivative of turmeric, is widely cultivated in Asia and is commonly used as a spice to add flavor and yellow, coloring to food. Curcumin is known to have anti-inflammatory and anti-oxidant effects in addition to acting as an anti-carcinogenic and neuroprotective agent [26–28]. Recent studies have reported a beneficial effect of curcumin on the cardiovascular system, suggesting its potential as a therapeutic agent for the treatment of cardiovascular disease and endothelial dysfunction [19]. In this study, we showed an improvement in endothelial function after 8 weeks of the curcumin ingestion in postmenopausal women. The magnitude of the improvement achieved by curcumin treatment was comparable to that obtained with exercise. Therefore, regular ingestion of curcumin could be a preventive measure against cardiovascular disease in postmenopausal women. Furthermore, our results suggest that curcumin may be a potential alternative treatment for patients who are unable to exercise.

Our hypothesis that curcumin ingestion may improve endothelial function in a similar manner as exercise was




Fig. 2 – Changes in FMD in response to intervention. Data are expressed as means \pm SD. * $P < .05$ vs control group.

proven in the present study, as shown in the significant and comparable increase in FMD after 8 weeks of curcumin ingestion and exercise training. Regular aerobic exercise improves enhanced endothelial function, which has been associated with increased nitric oxide bioavailability [9,29]. Therefore, the exercise-induced endothelial function improvement in our study could have been mediated by an increase in nitric oxide bioavailability. However, the mechanism responsible for the curcumin ingestion induced improvement in endothelial function is unclear. Curcumin exerts anti-inflammatory and anti-oxidative effects by inhibiting tumor necrosis factor- α (TNF- α) [30], suggesting that its effect on endothelial function may be mediated by the suppression of inflammation and/or oxidative stress via down-regulation of TNF- α . However, TNF- α levels were not assessed in this study. Further studies are warranted to clarify the mechanism underlying the effect of curcumin on endothelial function.

In our previous study, we investigated the effect of exercise training and curcumin ingestion on central arterial hemodynamics, that is, wave reflection and central blood pressure [21]. Wave reflection and central blood pressure did not improve with exercise training alone or curcumin ingestion alone, whereas these parameters improved in response to the combination of exercise training with curcumin ingestion. However, in the present study, we investigated the effect of curcumin on endothelial function as measured by FMD and demonstrated for the first time that curcumin ingestion alone increased FMD, similar to the effect of exercise demonstrated previously. The effect of the combination of exercise training with curcumin ingestion on vascular endothelial function were not assessed in this study and should be addressed in future studies in comparison to the effect of each agent alone.

The present study had some limitations. First, this study was conducted with a small sample size in each group. However, the power calculation indicated that the number of subjects was sufficient to establish statistical significance. Furthermore, this study focused only on postmenopausal women. Thus, the findings of this study cannot be generalized to other populations, such as men and young women. Second, the present study measured only FMD as an index of endothelial function. We did not measure any biomarkers of inflammation and oxidative stress, plasma curcumin concentration, or endothelial function using different indexes. Further studies are needed to clarify the mechanism underlying the effect of curcumin on vascular endothelial function. Third, analysis of the baseline characteristics of the participants and the baseline FMD suggest that the subjects in the exercise group showed overall better before the start of the study. Although there were no significant differences in any of the variables among the groups before the intervention, small differences in baseline values might have biased the results.

In conclusion, we examined the effect of curcumin ingestion and exercise training on endothelial function in postmenopausal women and found that regular curcumin ingestion and exercise improve endothelial function. These findings suggest that curcumin could be developed as a therapeutic strategy for the treatment of cardiovascular disease similar to exercise training in postmenopausal women.

Acknowledgment

This work was supported by Grants-in-Aid for Scientific Research 21300234 and 2160179 from Japan Society for the Promotion of Science.

REFERENCES

- [1] Moncada S, Palmer RM, Higgs EA. Nitric oxide: physiology, pathophysiology, and pharmacology. *Pharmacol Rev* 1991;43: 109–42.
- [2] Thijssen DH, Black MA, Pyke KE, Padilla J, Atkinson G, Harris RA, et al. Assessment of flow-mediated dilation in humans: a methodological and physiological guideline. *Am J Physiol Heart Circ Physiol* 2011;300:H2–H12.
- [3] Celermajer DS, Sorensen KE, Spiegelhalter DJ, Georgakopoulos D, Robinson J, Deanfield JE. Aging is associated with endothelial dysfunction in healthy men years before the age-related decline in women. *J Am Coll Cardiol* 1994;24:471–6.
- [4] Gates PE, Boucher ML, Silver AE, Monahan KD, Seals DR. Impaired flow-mediated dilation with age is not explained by L-arginine bioavailability or endothelial asymmetric dimethylarginine protein expression. *J Appl Physiol* 2007;102: 63–71.
- [5] Taddei S, Virdis A, Ghiadoni L, Mattei P, Sudano I, Bernini G, et al. Menopause is associated with endothelial dysfunction in women. *Hypertension* 1996;28:576–82.
- [6] Gavin KM, Seals DR, Silver AE, Moreau KL. Vascular endothelial estrogen receptor alpha is modulated by estrogen status and related to endothelial function and endothelial nitric oxide synthase in healthy women. *J Clin Endocrinol Metab* 2009;94:3513–20.
- [7] La Vecchia C. Sex hormones and cardiovascular risk. *Hum Reprod* 1992;7:162–7.
- [8] DeSouza CA, Shapiro LF, Clevenger CM, Dinunno FA, Monahan KD, Tanaka H, et al. Regular aerobic exercise prevents and restores age-related declines in endothelium-dependent vasodilation in healthy men. *Circulation* 2000;102: 1351–7.
- [9] Taddei S, Galetta F, Virdis A, Ghiadoni L, Salvetti G, Franzoni F, et al. Physical activity prevents age-related impairment in nitric oxide availability in elderly athletes. *Circulation* 2000;101:2896–901.
- [10] Yoshizawa M, Maeda S, Miyaki A, Misono M, Choi Y, Shimojo N, et al. Additive beneficial effects of lactotripeptides intake with regular exercise on endothelium-dependent dilatation in postmenopausal women. *Am J Hypertens* 2010;23:368–72.
- [11] Herrera MD, Mingorance C, Rodríguez-Rodríguez R, Alvarez de Sotomayor M. Endothelial dysfunction and aging: an update. *Ageing Res Rev* 2010;9:142–52.
- [12] Donato AJ, Black AD, Jablonski KL, Gano LB, Seals DR. Aging is associated with greater nuclear NF kappa B, reduced I kappa B alpha, and increased expression of proinflammatory cytokines in vascular endothelial cells of healthy humans. *Ageing Cell* 2008;7:805–12.
- [13] Csiszar A, Ungvari Z, Edwards JG, Kaminski P, Wolin MS, Koller A, et al. Aging-induced phenotypic changes and oxidative stress impair coronary arteriolar function. *Circ Res* 2002;90:1159–66.
- [14] Goel A, Kunnumakkara AB, Aggarwal BB. Curcumin as “Curcumin”: from kitchen to clinic. *Biochem Pharmacol* 2008;75:787–809.
- [15] Gupta SC, Patchva S, Koh W, Aggarwal BB. Discovery of curcumin, a component of golden spice, and its miraculous biological activities. *Clin Exp Pharmacol Physiol* 2012;39: 283–99.
- [16] Gupta SC, Prasad S, Kim JH, Patchva S, Webb LJ, Priyadarsini IK, et al. Multitargeting by curcumin as revealed by molecular interaction studies. *Nat Prod Rep* 2011;28:1937–55.
- [17] Cho JW, Lee KS, Kim GW. Curcumin attenuate the expression of IL-1 β , IL-6, and TNF- α as well as cyclin E in TNF- α -treated HaCaT cell; NF- κ B and MAPKs as potential upstream targets. *Int J Mol Med* 2007;19:469–74.
- [18] Quiles JL, Mesa MD, Ramírez-Tortosa CL, Aguilera CM, Battionio M, Gil A, et al. Curcuma longa extract supplementation reduces oxidative stress and attenuates aortic fatty streak development in rabbits. *Arterioscler Thromb Vasc Biol* 2002;22:1225–31.
- [19] Wongcharoen W, Phrommintikul A. The protective role of curcumin in cardiovascular diseases. *Int J Cardiol* 2009;133: 145–51.
- [20] Kanai M, Imaizumi A, Otsuka Y, Sasaki H, Hashiguchi M, Tsujiko K, et al. Dose-escalation and pharmacokinetic study of nanoparticle curcumin, a potential anticancer agent with improved bioavailability, in healthy human volunteers. *Cancer Chemother Pharmacol* 2012;69:65–70.
- [21] Sugawara J, Akazawa N, Miyaki A, Choi Y, Tanabe Y, Imai T, et al. Effect of endurance exercise training and curcumin intake on central arterial hemodynamics in postmenopausal women: pilot study. *Am J Hypertens* 2012;25:651–6.
- [22] Sasaki H, Sunagawa Y, Takahashi K, Imaizumi A, Fukuda H, Hashimoto T, et al. Innovative preparation of curcumin for improved oral bioavailability. *Biol Pharm Bull* 2011;34:660–5.
- [23] Yoshizawa M, Maeda S, Miyaki A, Misono M, Choi Y, Shimojo N, et al. Additive beneficial effects of lactotripeptides and aerobic exercise on arterial compliance in postmenopausal women. *Am J Physiol Heart Circ Physiol* 2009;297:H1899–903.
- [24] Kawano N, Emoto M, Mori K, Yamazaki Y, Urata H, Tsuchikura S, et al. Association of endothelial and vascular smooth muscle dysfunction with cardiovascular risk factors, vascular complications, and subclinical carotid atherosclerosis in type 2 diabetic patients. *J Atheroscler Thromb* 2011 [In press].
- [25] Miyaki A, Maeda S, Otsuki T, Ajisaka R. Plasma pentraxin 3 concentration increases in endurance-trained men. *Med Sci Sports Exerc* 2011;43:12–7.
- [26] Sharma RA, Gescher AJ, Steward WP. Curcumin: the story so far. *Eur J Cancer* 2005;41:1955–68.
- [27] Maheshwari RK, Singh AK, Gaddipati J, Srimal RC. Multiple biological activities of curcumin: a short review. *Life Sci* 2006;78:2081–7.
- [28] Jiang J, Wang W, Sun YJ, Hu M, Li F, Zhu DY. Neuroprotective effect of curcumin on focal cerebral ischemic rats by preventing blood-brain barrier damage. *Eur J Pharmacol* 2007;561:54–62.
- [29] Eskurza I, Monahan KD, Robinson JA, Seals DR. Effect of acute and chronic ascorbic acid on flow-mediated dilatation with sedentary and physically active human ageing. *J Physiol* 2004;556:315–24.
- [30] He ZY, Shi CB, Wen H, Li FL, Wang BL, Wang J. Upregulation of p53 expression in patients with colorectal cancer by administration of curcumin. *Cancer Invest* 2011;29:208–13.